

## Free $n$ -dinilpotent doppelsemigroups

Anatolii V. Zhuchok\* and Milan Demko\*\*

**ABSTRACT.** A doppelalgebra is an algebra defined on a vector space with two binary linear associative operations. Doppelalgebras play a prominent role in algebraic  $K$ -theory. In this paper we consider doppelsemigroups, that is, sets with two binary associative operations satisfying the axioms of a doppelalgebra. We construct a free  $n$ -dinilpotent doppelsemigroup and study separately free  $n$ -dinilpotent doppelsemigroups of rank 1. Moreover, we characterize the least  $n$ -dinilpotent congruence on a free doppelsemigroup, establish that the semigroups of the free  $n$ -dinilpotent doppelsemigroup are isomorphic and the automorphism group of the free  $n$ -dinilpotent doppelsemigroup is isomorphic to the symmetric group. We also give different examples of doppelsemigroups and prove that a system of axioms of a doppelsemigroup is independent.

### 1. Introduction

The notion of a doppelalgebra was considered by Richter [1] in the context of algebraic  $K$ -theory. She defined this notion as a vector space over a field equipped with two binary linear associative operations  $\dashv$  and  $\vdash$  satisfying the axioms  $(x \dashv y) \vdash z = x \dashv (y \vdash z)$ ,  $(x \vdash y) \dashv z = x \vdash (y \dashv z)$ . Observe that any doppelalgebra gives rise to a Lie algebra by  $[x, y] = x \vdash y + x \dashv y - y \vdash x - y \dashv x$  and conversely, any

---

\*The paper was written during the research stay of the first author at the University of Presov under the National Scholarship Programme of the Slovak Republic.

\*\*The second author acknowledges the support of the Slovak VEGA Grant No. 1/0063/14.

**2010 MSC:** 08B20, 20M10, 20M50, 17A30.

**Key words and phrases:** doppelalgebra, interassociativity, doppelsemigroup, free  $n$ -dinilpotent doppelsemigroup, free doppelsemigroup, semigroup, congruence.

Lie algebra has a universal enveloping doppelalgebra (see [1]). Moreover, for any doppelalgebra a new operation  $\cdot$  defined by  $x \cdot y = x \vdash y + x \dashv y$  is associative and so, there exists a functor from the category of doppelalgebras to the category of associative algebras. Later Pirashvili [2] considered duplexes which are sets equipped with two binary associative operations and constructed a free duplex of an arbitrary rank. He also considered duplexes with operations satisfying the axioms of a doppelalgebra denoting obtained category by  $\text{Duplexes}_2$ . Such algebraic structures are called doppelsemigroups [3]. A free doppelsemigroup of rank 1 is given in [1] (see also [2]). Operations of the free doppelsemigroup of rank 1 are used in [4]. Doppelalgebras appeared in [5] as algebras over some operad. Doppelalgebras and doppelsemigroups have relationships with interassociativity for semigroups originated by Drouzy [6] and investigated in [7–11], strong interassociativity for semigroups introduced by Gould and Richardson [12] and dimonoids introduced by Loday [13] (see also [14–18], [20–22]). Doppelsemigroups are a generalization of semigroups and all results obtained for doppelsemigroups can be applied to doppelalgebras. For further details and background see [1].

The free product of doppelsemigroups, the free doppelsemigroup, the free commutative doppelsemigroup and the free  $n$ -nilpotent doppelsemigroup were constructed in [3]. The paper [17] gives a classification of relatively free dimonoids, in particular, therein the free  $n$ -dinilpotent dimonoid [18] is presented. In this paper we continue researches from [3, 18] developing the variety theory of doppelsemigroups. The main focus of our paper is to study dinilpotent doppelsemigroups.

In Section 3 we present different examples of doppelsemigroups.

In Section 4 we prove that a system of axioms of a doppelsemigroup is independent.

In Section 5 we construct a free  $n$ -dinilpotent doppelsemigroup of an arbitrary rank and consider separately free  $n$ -dinilpotent doppelsemigroups of rank 1. We also establish that the semigroups of the free  $n$ -dinilpotent doppelsemigroup are isomorphic and the automorphism group of the free  $n$ -dinilpotent doppelsemigroup is isomorphic to the symmetric group.

In the final section we characterize the least  $n$ -dinilpotent congruence on a free doppelsemigroup.

## 2. Preliminaries

Recall that a doppelalgebra [1, 2] is a vector space  $V$  over a field equipped with two binary linear operations  $\dashv$  and  $\vdash: V \otimes V \rightarrow V$ , satisfying

the axioms

$$(x \dashv y) \vdash z = x \dashv (y \vdash z), \quad (\text{D1})$$

$$(x \vdash y) \dashv z = x \vdash (y \dashv z), \quad (\text{D2})$$

$$(x \dashv y) \dashv z = x \dashv (y \dashv z), \quad (\text{D3})$$

$$(x \vdash y) \vdash z = x \vdash (y \vdash z). \quad (\text{D4})$$

A nonempty set with two binary operations  $\dashv$  and  $\vdash$  satisfying the axioms (D1)–(D4) is called a doppelsemigroup [3].

Given a semigroup  $(D, \dashv)$ , consider a semigroup  $(D, \vdash)$  defined on the same set. Recall that  $(D, \vdash)$  is an interassociate of  $(D, \dashv)$  [6], if the axioms (D1) and (D2) hold. Strong interassociativity [12] is defined by the axioms (D1) and (D2) along with

$$x \vdash (y \dashv z) = x \dashv (y \vdash z). \quad (\text{D5})$$

Thus, we can see that in any doppelsemigroup  $(D, \dashv, \vdash)$ ,  $(D, \vdash)$  is an interassociate of  $(D, \dashv)$ , and conversely, if a semigroup  $(D, \vdash)$  is an interassociate of a semigroup  $(D, \dashv)$ , then  $(D, \dashv, \vdash)$  is a doppelsemigroup [3]. Moreover, a semigroup  $(D, \vdash)$  is a strong interassociate of a semigroup  $(D, \dashv)$  if and only if  $(D, \dashv, \vdash)$  is a doppelsemigroup satisfying the axiom (D5).

Descriptions of all interassociates of a monogenic semigroup and of the free commutative semigroup are presented in [7] and [8, 10], respectively. More recently, the paper [11] was devoted to studying interassociates of the bicyclic semigroup. Methods of constructing interassociates for semigroups were developed in [19].

Recall the definition of a  $k$ -nilpotent semigroup (see also [14, 17, 18]). As usual,  $\mathbb{N}$  denotes the set of all positive integers. A semigroup  $S$  is called nilpotent, if  $S^{n+1} = 0$  for some  $n \in \mathbb{N}$ . The least such  $n$  is called the nilpotency index of  $S$ . For  $k \in \mathbb{N}$  a nilpotent semigroup of nilpotency index  $\leq k$  is called  $k$ -nilpotent.

An element  $0$  of a doppelsemigroup  $(D, \dashv, \vdash)$  is called zero [3], if  $x * 0 = 0 = 0 * x$  for all  $x \in D$  and  $*$   $\in \{\dashv, \vdash\}$ . A doppelsemigroup  $(D, \dashv, \vdash)$  with zero will be called dinilpotent, if  $(D, \dashv)$  and  $(D, \vdash)$  are nilpotent semigroups. A dinilpotent doppelsemigroup  $(D, \dashv, \vdash)$  will be called  $n$ -dinilpotent, if  $(D, \dashv)$  and  $(D, \vdash)$  are  $n$ -nilpotent semigroups. If  $\rho$  is a congruence on a doppelsemigroup  $(D, \dashv, \vdash)$  such that  $(D, \dashv, \vdash)/\rho$  is an  $n$ -dinilpotent doppelsemigroup, we say that  $\rho$  is an  $n$ -dinilpotent congruence.

Note that operations of any 1-dinilpotent doppelsemigroup coincide and it is a zero semigroup. The class of all  $n$ -dinilpotent doppelsemigroups forms a subvariety of the variety of doppelsemigroups. It is not difficult to check that the variety of  $n$ -nilpotent doppelsemigroups [3] is a subvariety of the variety of  $n$ -dinilpotent doppelsemigroups. A doppelsemigroup which is free in the variety of  $n$ -dinilpotent doppelsemigroups will be called a free  $n$ -dinilpotent doppelsemigroup.

**Lemma 1** ([3], Lemma 3.1). *In a doppelsemigroup  $(D, \dashv, \vdash)$  for any  $n > 1, n \in \mathbb{N}$ , and any  $x_i \in D, 1 \leq i \leq n + 1$ , and  $*_j \in \{\dashv, \vdash\}, 1 \leq j \leq n$ , any parenthesizing of*

$$x_1 *_1 x_2 *_2 \dots *_n x_{n+1}$$

*gives the same element from  $D$ .*

The free doppelsemigroup is given in [3]. Recall this construction.

Let  $X$  be an arbitrary nonempty set and let  $\omega$  be an arbitrary word in the alphabet  $X$ . The length of  $\omega$  will be denoted by  $l_\omega$ . Let further  $F[X]$  be the free semigroup on  $X$ ,  $T$  the free monoid on the two-element set  $\{a, b\}$  and  $\theta \in T$  the empty word. By definition, the length  $l_\theta$  of  $\theta$  is equal to 0. Define operations  $\dashv$  and  $\vdash$  on  $F = \{(w, u) \in F[X] \times T \mid l_w - l_u = 1\}$  by

$$\begin{aligned} (w_1, u_1) \dashv (w_2, u_2) &= (w_1 w_2, u_1 a u_2), \\ (w_1, u_1) \vdash (w_2, u_2) &= (w_1 w_2, u_1 b u_2) \end{aligned}$$

for all  $(w_1, u_1), (w_2, u_2) \in F$ . The algebra  $(F, \dashv, \vdash)$  is denoted by  $\text{FDS}(X)$ .

**Theorem 1** ([3], Theorem 3.5).  *$\text{FDS}(X)$  is the free doppelsemigroup.*

If  $f : D_1 \rightarrow D_2$  is a homomorphism of doppelsemigroups, the corresponding congruence on  $D_1$  will be denoted by  $\Delta_f$ . Denote the symmetric group on  $X$  by  $\mathfrak{S}[X]$  and the automorphism group of a doppelsemigroup  $M$  by  $\text{Aut } M$ .

### 3. Some examples

In this section we give different examples of doppelsemigroups.

- a) Every semigroup can be considered as a doppelsemigroup (see [3]).
- b) Recall that a dimonoid [13–18, 20–22] is a nonempty set equipped with two binary operations  $\dashv$  and  $\vdash$  satisfying the axioms (D2)–(D4) and

$$\begin{aligned} (x \dashv y) \dashv z &= x \dashv (y \vdash z), \\ (x \dashv y) \vdash z &= x \vdash (y \vdash z). \end{aligned}$$

A dimonoid is called commutative [20], if both its operations are commutative. The following assertion gives relationships between commutative dimonoids and doppelsemigroups (this assertion was formulated without the proof in [3] and [16]).

**Proposition 1.** *Every commutative dimonoid is a doppelsemigroup.*

*Proof.* Let  $(D, \dashv, \vdash)$  be a commutative dimonoid. Then, by definition,  $(D, \dashv, \vdash)$  satisfies the axioms (D2)–(D4). From Lemma 2 of [20] it follows that  $(D, \dashv, \vdash)$  satisfies the axiom (D1). So, it is a doppelsemigroup.  $\square$

Examples of commutative dimonoids can be found in [14, 20].

c) Let  $(D, \dashv, \vdash)$  be a doppelsemigroup and  $a, b \in D$ . Define operations  $\dashv_a$  and  $\vdash_b$  on  $D$  by

$$x \dashv_a y = x \dashv a \dashv y, \quad x \vdash_b y = x \vdash b \vdash y$$

for all  $x, y \in D$ . By a direct verification  $(D, \dashv_a, \vdash_b)$  is a doppelsemigroup. We call the doppelsemigroup  $(D, \dashv_a, \vdash_b)$  a variant of  $(D, \dashv, \vdash)$ , or, alternatively, the sandwich doppelsemigroup of  $(D, \dashv, \vdash)$  with respect to the sandwich elements  $a$  and  $b$ , or the doppelsemigroup with deformed multiplications.

d) The direct product  $\prod_{i \in I} D_i$  of doppelsemigroups  $D_i, i \in I$ , is, obviously, a doppelsemigroup.

e) Now we give a new class of doppelsemigroups with zero.

Let  $\overline{D} = (D, \dashv, \vdash)$  be an arbitrary doppelsemigroup and  $I$  an arbitrary nonempty set. Define operations  $\dashv'$  and  $\vdash'$  on  $D' = (I \times D \times I) \cup \{0\}$  by

$$(i, a, j) *' (k, b, t) = \begin{cases} (i, a * b, t), & j = k, \\ 0, & j \neq k, \end{cases}$$

$$(i, a, j) *' 0 = 0 *' (i, a, j) = 0 *' 0 = 0$$

for all  $(i, a, j), (k, b, t) \in D' \setminus \{0\}$  and  $* \in \{\dashv, \vdash\}$ . The algebra  $(D', \dashv', \vdash')$  will be denoted by  $B(\overline{D}, I)$ .

**Proposition 2.**  *$B(\overline{D}, I)$  is a doppelsemigroup with zero.*

*Proof.* The proof is similar to the proof of Proposition 1 from [21].  $\square$

Observe that if operations of a doppelsemigroup  $\overline{D}$  coincide and it is a group  $G$ , then any Brandt semigroup [23] is isomorphic to some semigroup  $B(G, I)$ . So,  $B(\overline{D}, I)$  generalizes the semigroup  $B(G, I)$ . We call the doppelsemigroup  $B(\overline{D}, I)$  a Brandt doppelsemigroup.

#### 4. Independence of axioms of a doppelsemigroup

In this section for a doppelsemigroup we prove the following theorem.

**Theorem 2.** *A system of axioms (D1)–(D4) as defined above is independent.*

*Proof.* Let  $X$  be an arbitrary nonempty set,  $|X| > 1$ . Define operations  $\dashv$  and  $\vdash$  on  $X$  by

$$x \dashv y = x, \quad x \vdash y = y$$

for all  $x, y \in X$ . The model  $(X, \dashv, \vdash)$  satisfies the axioms (D2)–(D4) but does not satisfy (D1). Indeed, for all  $x, y, z \in X$ ,

$$\begin{aligned} (x \vdash y) \dashv z &= y = x \vdash (y \dashv z), \\ (x \dashv y) \dashv z &= x = x \dashv (y \dashv z), \\ (x \vdash y) \vdash z &= z = x \vdash (y \vdash z). \end{aligned}$$

Since  $|X| > 1$ , there is  $x, z \in X$  such that  $x \neq z$ . Consequently, for all  $y \in X$ ,

$$(x \dashv y) \vdash z = z \neq x = x \dashv (y \vdash z).$$

Put

$$x \dashv y = y, \quad x \vdash y = x$$

for all  $x, y \in X$ . As in the previous case, we can show that  $(X, \dashv, \vdash)$  satisfies the axioms (D1), (D3), (D4) but does not satisfy (D2).

Let  $\mathbb{N}^0$  be the set of all positive integers with zero and let

$$x \dashv y = 2x, \quad z \dashv 0 = 0 = 0 \dashv z, \quad z \vdash c = 0$$

for all  $x, y \in \mathbb{N}$  and  $z, c \in \mathbb{N}^0$ . In this case the model  $(\mathbb{N}^0, \dashv, \vdash)$  satisfies the axioms (D1), (D2), (D4) but does not satisfy (D3). Indeed, for all  $z, c, a \in \mathbb{N}^0$ ,

$$\begin{aligned} (z \dashv c) \vdash a &= 0 = z \dashv (c \vdash a), \\ (z \vdash c) \dashv a &= 0 = z \vdash (c \dashv a), \\ (z \vdash c) \vdash a &= 0 = z \vdash (c \vdash a). \end{aligned}$$

In addition, for all  $x, y, b \in \mathbb{N}$  we get

$$(x \dashv y) \vdash b = 2x \dashv b = 4x \neq 2x = x \dashv 2y = x \dashv (y \vdash b).$$

Put

$$z \dashv c = 0, \quad x \vdash y = 2y, \quad z \vdash 0 = 0 = 0 \vdash z$$

for all  $z, c \in \mathbb{N}^0$  and  $x, y \in \mathbb{N}$ . As in the previous case, we can show that  $(\mathbb{N}^0, \dashv, \vdash)$  satisfies the axioms (D1)–(D3) but does not satisfy (D4).  $\square$

### 5. Constructions

In this section we construct a free  $n$ -dinilpotent doppelsemigroup of an arbitrary rank and consider separately free  $n$ -dinilpotent doppelsemigroups of rank 1. We also establish that the semigroups of the free  $n$ -dinilpotent doppelsemigroup are isomorphic and the automorphism group of the free  $n$ -dinilpotent doppelsemigroup is isomorphic to the symmetric group.

As in Section 2, let  $F[X]$  be the free semigroup on  $X$ ,  $T$  the free monoid on the two-element set  $\{a, b\}$  and  $\theta \in T$  the empty word. For  $x \in \{a, b\}$  and all  $u \in T$ , the number of occurrences of an element  $x$  in  $u$  is denoted by  $d_x(u)$ . Obviously,  $d_x(\theta) = 0$ . Fix  $n \in \mathbb{N}$  and assume

$$M_n = \{(w, u) \in F[X] \times T \mid l_w - l_u = 1, d_x(u) + 1 \leq n, x \in \{a, b\}\} \cup \{0\}.$$

Define operations  $\dashv$  and  $\vdash$  on  $M_n$  by

$$(w_1, u_1) \dashv (w_2, u_2) = \begin{cases} (w_1w_2, u_1au_2), & d_x(u_1au_2) + 1 \leq n, x \in \{a, b\}, \\ 0, & \text{in all other cases,} \end{cases}$$

$$(w_1, u_1) \vdash (w_2, u_2) = \begin{cases} (w_1w_2, u_1bu_2), & d_x(u_1bu_2) + 1 \leq n, x \in \{a, b\}, \\ 0, & \text{in all other cases,} \end{cases}$$

$$(w_1, u_1) * 0 = 0 * (w_1, u_1) = 0 * 0 = 0$$

for all  $(w_1, u_1), (w_2, u_2) \in M_n \setminus \{0\}$  and  $* \in \{\dashv, \vdash\}$ . The obtained algebra will be denoted by  $\text{FDDS}_n(X)$ .

**Theorem 3.**  $\text{FDDS}_n(X)$  is the free  $n$ -dinilpotent doppelsemigroup.

*Proof.* First prove that  $\text{FDDS}_n(X)$  is a doppelsemigroup. Let  $(w_1, u_1), (w_2, u_2), (w_3, u_3) \in M_n \setminus \{0\}$ . For  $x, y, z \in \{a, b\}$  it is clear that

$$d_x(u_1yu_2zu_3) + 1 \leq n$$

implies

$$d_x(u_1yu_2) + 1 \leq n, \tag{1}$$

$$d_x(u_2zu_3) + 1 \leq n. \tag{2}$$

Let  $d_x(u_1au_2au_3) + 1 \leq n$  for all  $x \in \{a, b\}$ . Then, using (1), (2), we get

$$\begin{aligned} ((w_1, u_1) \dashv (w_2, u_2)) \dashv (w_3, u_3) &= (w_1w_2, u_1au_2) \dashv (w_3, u_3) \\ &= (w_1w_2w_3, u_1au_2au_3) \\ &= (w_1, u_1) \dashv (w_2w_3, u_2au_3) \\ &= (w_1, u_1) \dashv ((w_2, u_2) \dashv (w_3, u_3)). \end{aligned}$$

If  $d_x(u_1au_2au_3) + 1 > n$  for some  $x \in \{a, b\}$ , then, obviously,

$$((w_1, u_1) \dashv (w_2, u_2)) \dashv (w_3, u_3) = 0 = (w_1, u_1) \dashv ((w_2, u_2) \dashv (w_3, u_3)).$$

So, the axiom (D3) of a doppelsemigroup holds.

If  $d_x(u_1au_2bu_3) + 1 \leq n$  for all  $x \in \{a, b\}$ , then, using (1), (2), obtain

$$\begin{aligned} ((w_1, u_1) \dashv (w_2, u_2)) \vdash (w_3, u_3) &= (w_1w_2, u_1au_2) \vdash (w_3, u_3) \\ &= (w_1w_2w_3, u_1au_2bu_3) \\ &= (w_1, u_1) \dashv (w_2w_3, u_2bu_3) \\ &= (w_1, u_1) \dashv ((w_2, u_2) \vdash (w_3, u_3)). \end{aligned}$$

Let  $d_x(u_1au_2bu_3) + 1 > n$  for some  $x \in \{a, b\}$ . Then, clearly,

$$((w_1, u_1) \dashv (w_2, u_2)) \vdash (w_3, u_3) = 0 = (w_1, u_1) \dashv ((w_2, u_2) \vdash (w_3, u_3)).$$

Thus, the axiom (D1) of a doppelsemigroup holds. Similarly, one can check the axioms (D2) and (D4). Thus,  $\text{FDDS}_n(X)$  is a doppelsemigroup.

Take arbitrary elements  $(w_i, u_i) \in M_n \setminus \{0\}$ ,  $1 \leq i \leq n+1$ . It is clear that

$$d_a(u_1au_2a \dots au_{n+1}) + 1 > n.$$

From here

$$(w_1, u_1) \dashv (w_2, u_2) \dashv \dots \dashv (w_{n+1}, u_{n+1}) = 0.$$

At the same time, assuming  $y^0 = \theta$  for  $y \in \{a, b\}$ , for any  $(x_i, \theta) \in M_n \setminus \{0\}$ , where  $x_i \in X$ ,  $1 \leq i \leq n$ , get

$$(x_1, \theta) \dashv (x_2, \theta) \dashv \dots \dashv (x_n, \theta) = (x_1x_2 \dots x_n, a^{n-1}) \neq 0.$$

From the last arguments we conclude that  $(M_n, \dashv)$  is a nilpotent semigroup of nilpotency index  $n$ . Analogously, we can prove that  $(M_n, \vdash)$  is a nilpotent semigroup of nilpotency index  $n$ . So, by definition,  $\text{FDDS}_n(X)$  is an  $n$ -dinilpotent doppelsemigroup.

Let us show that  $\text{FDDS}_n(X)$  is free in the variety of  $n$ -dinilpotent doppelsemigroups.

Obviously,  $\text{FDDS}_n(X)$  is generated by  $X \times \{\theta\}$ . Let  $(K, \dashv', \vdash')$  be an arbitrary  $n$ -dinilpotent doppelsemigroup. Let  $\beta : X \times \{\theta\} \rightarrow K$  be an arbitrary map. Consider a map  $\alpha : X \rightarrow K$  such that  $x\alpha = (x, \theta)\beta$  for all  $x \in X$  and define a map

$$\pi : \text{FDDS}_n(X) \rightarrow (K, \dashv', \vdash')$$



by

$$\omega\pi = \begin{cases} x_1\alpha\tilde{y}_1x_2\alpha\tilde{y}_2\dots\tilde{y}_{s-1}x_s\alpha, & \text{if } \omega = (x_1x_2\dots x_s, y_1y_2\dots y_{s-1}), \\ & x_d \in X, 1 \leq d \leq s, y_p \in \{a, b\}, \\ & 1 \leq p \leq s-1, s > 1, \\ x_1\alpha, & \text{if } \omega = (x_1, \theta), x_1 \in X, \\ 0, & \text{if } \omega = 0, \end{cases}$$

where

$$\tilde{y}_p = \begin{cases} \dashv', & y_p = a, \\ \vdash', & y_p = b \end{cases}$$

for all  $1 \leq p \leq s-1, s > 1$ . According to Lemma 1  $\pi$  is well-defined.

To show that  $\pi$  is a homomorphism we will use the axioms of a doppelsemigroup and the identities of an  $n$ -dinilpotent doppelsemigroup.

If  $s = 1$ , we will regard the sequence  $y_1y_2\dots y_{s-1} \in T$  as  $\theta$ . For arbitrary elements

$$\begin{aligned} (w_1, u_1) &= (x_1x_2\dots x_s, y_1y_2\dots y_{s-1}), \\ (w_2, u_2) &= (z_1z_2\dots z_k, c_1c_2\dots c_{k-1}) \in \text{FDDS}_n(X), \end{aligned}$$

where  $x_d, z_i \in X, 1 \leq d \leq s, 1 \leq i \leq k, y_p, c_j \in \{a, b\}, 1 \leq p \leq s-1, 1 \leq j \leq k-1$ , in the case  $d_x(u_1au_2) + 1 \leq n$  for all  $x \in \{a, b\}$ , we get

$$\begin{aligned} &((x_1x_2\dots x_s, y_1y_2\dots y_{s-1}) \dashv (z_1z_2\dots z_k, c_1c_2\dots c_{k-1}))\pi \\ &= (x_1\dots x_s z_1\dots z_k, y_1\dots y_{s-1} a c_1\dots c_{k-1})\pi \\ &= x_1\alpha\tilde{y}_1\dots\tilde{y}_{s-1}x_s\alpha\tilde{a}z_1\alpha\tilde{c}_1\dots\tilde{c}_{k-1}z_k\alpha \\ &= (x_1\alpha\tilde{y}_1\dots\tilde{y}_{s-1}x_s\alpha) \dashv' (z_1\alpha\tilde{c}_1\dots\tilde{c}_{k-1}z_k\alpha) \\ &= (x_1x_2\dots x_s, y_1y_2\dots y_{s-1})\pi \dashv' (z_1z_2\dots z_k, c_1c_2\dots c_{k-1})\pi. \end{aligned}$$

If  $d_x(u_1au_2) + 1 > n$  for some  $x \in \{a, b\}$ , then

$$((x_1x_2\dots x_s, y_1y_2\dots y_{s-1}) \dashv (z_1z_2\dots z_k, c_1c_2\dots c_{k-1}))\pi = 0\pi = 0.$$

Since  $(K, \dashv', \vdash')$  is  $n$ -dinilpotent, we have

$$\begin{aligned} 0 &= x_1\alpha\tilde{y}_1\dots\tilde{y}_{s-1}x_s\alpha\tilde{a}z_1\alpha\tilde{c}_1\dots\tilde{c}_{k-1}z_k\alpha \\ &= (x_1\alpha\tilde{y}_1\dots\tilde{y}_{s-1}x_s\alpha) \dashv' (z_1\alpha\tilde{c}_1\dots\tilde{c}_{k-1}z_k\alpha) \\ &= (x_1x_2\dots x_s, y_1y_2\dots y_{s-1})\pi \dashv' (z_1z_2\dots z_k, c_1c_2\dots c_{k-1})\pi. \end{aligned}$$

So,

$$((w_1, u_1) \dashv (w_2, u_2))\pi = (w_1, u_1)\pi \dashv' (w_2, u_2)\pi$$

for all  $(w_1, u_1), (w_2, u_2) \in \text{FDDS}_n(X)$ .

Similarly for  $\vdash$ . So,  $\pi$  is a homomorphism. Clearly,  $(x, \theta)\pi = (x, \theta)\beta$  for all  $(x, \theta) \in X \times \{\theta\}$ . Since  $X \times \{\theta\}$  generates  $\text{FDDS}_n(X)$ , the uniqueness of such homomorphism  $\pi$  is obvious. Thus,  $\text{FDDS}_n(X)$  is free in the variety of  $n$ -dinilpotent doppelsemigroups.  $\square$

Now we construct a doppelsemigroup which is isomorphic to the free  $n$ -dinilpotent doppelsemigroup of rank 1.

Fix  $n \in \mathbb{N}$  and assume

$$\bar{\Phi}_n = \{u \in T \mid d_x(u) + 1 \leq n, x \in \{a, b\}\} \cup \{0\}.$$

Define operations  $\dashv$  and  $\vdash$  on  $\bar{\Phi}_n$  by

$$u_1 \dashv u_2 = \begin{cases} u_1 a u_2, & d_x(u_1 a u_2) + 1 \leq n, x \in \{a, b\}, \\ 0, & \text{in all other cases,} \end{cases}$$

$$u_1 \vdash u_2 = \begin{cases} u_1 b u_2, & d_x(u_1 b u_2) + 1 \leq n, x \in \{a, b\}, \\ 0, & \text{in all other cases,} \end{cases}$$

$$u_1 * 0 = 0 * u_1 = 0 * 0 = 0$$

for all  $u_1, u_2 \in \bar{\Phi}_n \setminus \{0\}$  and  $*$   $\in$   $\{\dashv, \vdash\}$ . The obtained algebra will be denoted by  $\Phi_n$ . Obviously,  $\Phi_n$  is a doppelsemigroup.

**Lemma 2.** *If  $|X| = 1$ , then  $\Phi_n \cong \text{FDDS}_n(X)$ .*

*Proof.* Let  $X = \{r\}$ . One can show that a map  $\gamma : \Phi_n \rightarrow \text{FDDS}_n(X)$ , defined by the rule

$$u\gamma = \begin{cases} (r^{l_u+1}, u), & u \in \bar{\Phi}_n \setminus \{0\}, \\ 0, & u = 0, \end{cases}$$

is an isomorphism.  $\square$

The following lemma establishes a relationship between semigroups of the free  $n$ -dinilpotent doppelsemigroup  $\text{FDDS}_n(X)$ .

**Lemma 3.** *The semigroups  $(M_n, \dashv)$  and  $(M_n, \vdash)$  are isomorphic.*

*Proof.* Let  $\widehat{a} = b$ ,  $\widehat{b} = a$  and define a map  $\sigma : (M_n, \dashv) \rightarrow (M_n, \vdash)$  by putting

$$t\sigma = \begin{cases} (w, \widehat{y_1}\widehat{y_2}\dots\widehat{y_m}), & t = (w, y_1y_2\dots y_m) \in M_n \setminus \{0\}, \\ & y_p \in \{a, b\}, 1 \leq p \leq m, \\ t, & \text{in all other cases.} \end{cases}$$

An immediate verification shows that  $\sigma$  is an isomorphism.  $\square$

Since the set  $X \times \{\theta\}$  is generating for  $\text{FDDS}_n(X)$ , we obtain the following description of the automorphism group of the free  $n$ -dinilpotent doppelsemigroup.

**Lemma 4.**  $\text{Aut FDDS}_n(X) \cong \mathfrak{S}[X]$ .

## 6. The least $n$ -dinilpotent congruence on a free doppelsemigroup

In this section we present the least  $n$ -dinilpotent congruence on a free doppelsemigroup.

Let  $\text{FDS}(X)$  be the free doppelsemigroup (see Section 2) and  $n \in \mathbb{N}$ . Define a relation  $\mu_{(n)}$  on  $\text{FDS}(X)$  by

$$(w_1, u_1)\mu_{(n)}(w_2, u_2) \quad \text{if and only if} \quad (w_1, u_1) = (w_2, u_2) \text{ or} \\ \begin{cases} d_x(u_1) + 1 > n & \text{for some } x \in \{a, b\}, \\ d_y(u_2) + 1 > n & \text{for some } y \in \{a, b\}. \end{cases}$$

**Theorem 4.** *The relation  $\mu_{(n)}$  on the free doppelsemigroup  $\text{FDS}(X)$  is the least  $n$ -dinilpotent congruence.*

*Proof.* Define a map  $\varphi : \text{FDS}(X) \rightarrow \text{FDDS}_n(X)$  by

$$(w, u)\varphi = \begin{cases} (w, u), & \text{if } d_x(u) + 1 \leq n \text{ for all } x \in \{a, b\}, \\ 0, & \text{in all other cases} \end{cases}$$

$((w, u) \in \text{FDS}(X))$ . Show that  $\varphi$  is a homomorphism.

Let  $(w_1, u_1), (w_2, u_2) \in \text{FDS}(X)$  and  $d_x(u_1au_2) + 1 \leq n$  for all  $x \in \{a, b\}$ . From the last inequality it follows that  $d_x(u_1) + 1 \leq n$  and  $d_x(u_2) + 1 \leq n$  for all  $x \in \{a, b\}$ . Then

$$\begin{aligned} ((w_1, u_1) \dashv (w_2, u_2))\varphi &= (w_1w_2, u_1au_2)\varphi = (w_1w_2, u_1au_2) \\ &= (w_1, u_1) \dashv (w_2, u_2) = (w_1, u_1)\varphi \dashv (w_2, u_2)\varphi. \end{aligned}$$

If  $d_x(u_1au_2) + 1 > n$  for some  $x \in \{a, b\}$ , then

$$((w_1, u_1) \dashv (w_2, u_2))\varphi = (w_1w_2, u_1au_2)\varphi = 0 = (w_1, u_1)\varphi \dashv (w_2, u_2)\varphi.$$

Let further  $d_x(u_1bu_2) + 1 \leq n$  for all  $x \in \{a, b\}$ . Then  $d_x(u_1) + 1 \leq n$ ,  $d_x(u_2) + 1 \leq n$  for all  $x \in \{a, b\}$  and

$$\begin{aligned} ((w_1, u_1) \vdash (w_2, u_2))\varphi &= (w_1w_2, u_1bu_2)\varphi = (w_1w_2, u_1bu_2) \\ &= (w_1, u_1) \vdash (w_2, u_2) = (w_1, u_1)\varphi \vdash (w_2, u_2)\varphi. \end{aligned}$$

If  $d_x(u_1bu_2) + 1 > n$  for some  $x \in \{a, b\}$ , then

$$((w_1, u_1) \vdash (w_2, u_2))\varphi = (w_1w_2, u_1bu_2)\varphi = 0 = (w_1, u_1)\varphi \vdash (w_2, u_2)\varphi.$$

Thus,  $\varphi$  is a surjective homomorphism. By Theorem 3  $\text{FDDS}_n(X)$  is the free  $n$ -dinilpotent doppelsemigroup. Then  $\Delta_\varphi$  is the least  $n$ -dinilpotent congruence on  $\text{FDS}(X)$ . From the definition of  $\varphi$  it follows that  $\Delta_\varphi = \mu_{(n)}$ .  $\square$

## References

- [1] B. Richter, *Dialgebren, Doppelalgebren und ihre Homologie*, Diplomarbeit, Universität Bonn (1997). <http://www.math.uni-hamburg.de/home/richter/publications.html>.
- [2] T. Pirashvili, *Sets with two associative operations*, Cent. Eur. J. Math. **2** (2003), 169–183.
- [3] A.V. Zhuchok, *Free products of doppelsemigroups*, Algebra Univers. Accepted in 2016.
- [4] J.-L. Loday, M.O. Ronco, *Order structure on the algebra of permutations and of planar binary trees*, J. Algebraic Combin. **15** (2002), 253–270.
- [5] M. Aguiar, M. Livernet, *The associative operad and the weak order on the symmetric groups*, J. Homotopy Relat. Struct. **2** (2007), no. 1, 57–84.
- [6] M. Drouzy, *La structuration des ensembles de semigroupes d'ordre 2, 3 et 4 par la relation d'interassociativité*. Manuscript (1986).
- [7] M. Gould, K.A. Linton, A.W. Nelson, *Interassociates of monogenic semigroups*, Semigroup Forum **68** (2004), 186–201.
- [8] B.N. Givens, K. Linton, A. Rosin, L. Dishman, *Interassociates of the free commutative semigroup on  $n$  generators*, Semigroup Forum **74** (2007), 370–378.
- [9] J.B. Hickey, *Semigroups under a sandwich operation*, Proc. Edinburgh Math. Soc. **26** (1983), 371–382.
- [10] A.B. Gorbakov, *Interassociativity on a free commutative semigroup*, Sib. Math. J. **54** (2013), no. 3, 441–445.
- [11] B.N. Givens, A. Rosin, K. Linton, *Interassociates of the bicyclic semigroup*, Semigroup Forum (2016). doi: 10.1007/s00233-016-9794-9.

- [12] M. Gould, R.E. Richardson, *Translational hulls of polynomially related semigroups*, Czechoslovak Math. J. **33** (1983), no. 1, 95–100.
- [13] J.-L. Loday, *Dialgebras*, In: Dialgebras and related operads, Lecture Notes in Math. Springer, Berlin **1763** (2001), 7–66.
- [14] A.V. Zhuchok, *Elements of dimonoid theory*, Mathematics and its Applications. Proceedings of Institute of Mathematics of NAS of Ukraine, Kiev **98** (2014), 304 p. (in Ukrainian).
- [15] A.V. Zhuchok, *Dimonoids and bar-units*, Sib. Math. J. **56** (2015), no. 5, 827–840.
- [16] A.V. Zhuchok, Yul.V. Zhuchok, *Free left  $n$ -dinilpotent dimonoids*, Semigroup Forum **93** (2016), no. 1, 161–179. doi: 10.1007/s00233-015-9743-z.
- [17] A.V. Zhuchok, *Structure of relatively free dimonoids*, Commun. Algebra **45** (2017), no. 4, 1639–1656, to appear. doi: 10.1080/00927872.2016.1222404.
- [18] A.V. Zhuchok, *Free  $n$ -dinilpotent dimonoids*, Problems of Physics, Mathematics and Technics **17** (2013), no. 4, 43–46.
- [19] S.J. Boyd, M. Gould, A.W. Nelson, *Interassociativity of semigroups*, Proceedings of the Tennessee Topology Conference, Nashville, TN, USA, 1996. Singapore: World Scientific (1997), 33–51.
- [20] A.V. Zhuchok, *Commutative dimonoids*, Algebra and Discrete Math. **2** (2009), 116–127.
- [21] A.V. Zhuchok, *Free  $n$ -nilpotent dimonoids*, Algebra and Discrete Math. **16** (2013), no. 2, 299–310.
- [22] A.V. Zhuchok, A.B. Gorbatkov, *On the structure of dimonoids*, Semigroup Forum (2016). doi: 10.1007/s00233-016-9795-8.
- [23] A.H. Clifford, G.B. Preston, *The algebraic theory of semigroups*, American Mathematical Society V. **1**, **2** (1964), (1967).

#### CONTACT INFORMATION

**A. V. Zhuchok**

Department of Algebra and System Analysis,  
Luhansk Taras Shevchenko National University,  
Gogol square, 1, Starobilsk, 92703, Ukraine  
*E-Mail(s)*: zhuchok\_a@mail.ru

**M. Demko**

Department of Physics, Mathematics and Tech-  
niques, University of Presov, Slovakia, 17. novem-  
bra 1, Presov, 08116, Slovakia  
*E-Mail(s)*: milan.demko@unipo.sk

Received by the editors: 03.10.2016  
and in final form 30.11.2016.